Technical Data Sheet

InVivoMAb anti-mouse CD4



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific* Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0003-3
Clone: YTS 177
Isotype: Rat IgG2a

Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: Mouse Spleen Cells

Reported Applications: in vivo blockade of CD4+ T-cell responses

Western blot

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB_1107642

Molecular Weight: 150 kDa

Description

The YTS 177 monoclonal antibody reacts with the mouse CD4. The CD4 antigen is a 55 kDa cell surface type I membrane glycoprotein belonging to the immunoglobulin superfamily. CD4 acts as a co-receptor which in cooperation with the T cell receptor (TCR) interacts with class II MHC molecules displayed by antigen presenting cells (APC). CD4 is expressed by the majority of thymocytes, most helper T cells, a subset of NK-T cells and weakly by dendritic cells and macrophages. CD4 plays an important role in the development of T cells and is required for mature T cells to function optimally. The YTS 177 antibody has been shown to compete with clones GK1.5 and YTS 191 for CD4 binding. Additionally, the YTS 177 antibody has been reported to be non-depleting but binding does induce rapid internalization of CD4 on both CD4+ Foxp3- T cells and Foxp3+ regulatory T cells. Further, the YTS 177 clone has been shown to suppress or even prevent allograft rejection, allergic reactions and autoimmune responses.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Bio X Cell, LLC Page 1 of 2

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0003-3#tab_references or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2