

Technical Data Sheet

InVivoMAb anti-mouse LFA-1 α (CD11a)



bioxcell.com

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Website Link: <https://bioxcell.com/invivomab-anti-mouse-lfa-1-alpha-cd11a-be0005-1>

Product Information

Catalog Number: BE0005-1
Clone: FD441.8
Isotype: Rat IgG2b
Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Cloned CTL B18 cells
Reported Applications: *in vivo* LFA-1 neutralization
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: ≤ 1 EU/mg (≤ 0.001 EU/ μ g)
Determined by LAL assay
Purity: $\geq 95\%$
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_1107574](https://abnova.com/AB_1107574)
Molecular Weight: 150 kDa

Description

The FD441.8 monoclonal antibody reacts with mouse LFA-1 α (lymphocyte function-associated antigen 1 alpha) also known as integrin alpha L chain and CD11a. LFA-1 α and CD18 combine to form LFA-1, a 180 kDa glycoprotein and a member of the integrin family. LFA-1 is expressed on the surface of all leukocytes including lymphocytes, monocytes, macrophages, and granulocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50), and also functions in lymphocyte costimulatory signaling.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/invivomab-anti-mouse-lfa-1-alpha-cd11a-be0005-1?utm_source=cr9k1b#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © Bio X Cell, LLC