Technical Data Sheet

InVivoMAb anti-mouse LFA-1α (CD11a)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific* Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0006
Clone: M17/4
Isotype: Rat IgG2a, κ

Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: C57BL/6 mouse splenic secondary cytotoxic T cells

Reported Applications: in vivo LFA-1 neutralization

Flow cytometry

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2ΕU/mg (<0.002ΕU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB_1107578

Molecular Weight: 150 kDa

Description

The M17/4 monoclonal antibody reacts with mouse LFA-1 α (lymphocyte function-associated antigen 1 alpha) also known as integrin alpha L chain and CD11a. LFA-1 α and CD18 combine to form LFA-1, a 180 kDa glycoprotein and a member of the integrin family. LFA-1 is expressed on the surface of all leukocytes including lymphocytes, monocytes, macrophages, and granulocytes. LFA-1 plays a central role in leukocyte intercellular adhesion through interactions with its ligands, ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50), and also functions in lymphocyte costimulatory signaling.

Storage

Store at the stock concentration at 4°C. Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

Bio X Cell, LLC Page 1 of 2

experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0006#tab_references or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC @ 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2