# **Technical Data Sheet**

#### InVivoMAb anti-mouse/human CD11b



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

## Lot Specific Information

Lot Number: Lot Specific\*
Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

#### **Product Information**

 Catalog Number:
 BE0007

 Clone:
 M1/70

 Isotype:
 Rat IgG2b, к

**Recommended Isotype Control(s):** InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer

Immunogen: B10 mouse spleen cells enriched for T cells

**Reported Applications:** in vivo CD11b neutralization

ILC2 cell purification Flow cytometry

**Formulation:** PBS, pH 7.0

Contains no stabilizers or preservatives

**Endotoxin:** <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

**Production:** Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB\_1107582

Molecular Weight: 150 kDa

#### **Description**

The M1/70 monoclonal antibody reacts with mouse and human CD11b, a 170 kDa membrane glycoprotein also known as integrin alpha M (ITGAM). CD11b belongs to the integrin alpha family and is primarily expressed on granulocytes and monocytes/macrophages but also expressed on dendritic cells, NK cells, and subsets of T and B cells. CD11b and CD18 combine to form Mac-1. Mac-1 functions as a complement receptor as well as a receptor for fibrinogen, factor X, and ICAM1.

### **Storage**

Store at the stock concentration at  $4^{\circ}$ C. **Do not freeze.** 

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/faqs">https://bioxcell.com/faqs</a>.

#### **Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

Bio X Cell, LLC Page 1 of 2

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### **Application References**

For a complete list of references, visit <a href="https://bioxcell.com/catalogsearch/result/?q=BE0007#tab\_references">https://bioxcell.com/catalogsearch/result/?q=BE0007#tab\_references</a> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2