

Technical Data Sheet

InVivoMAb anti-human CD54 (ICAM-1)



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0020-2
Clone: R6-5-D6
Isotype: Mouse IgG2a
Recommended Isotype Control(s): InVivoMAb mouse IgG2a isotype control, unknown specificity
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: EBV transformed lymphoblast cell line
Reported Applications: *in vitro* T cell stimulation/activation
Immunofluorescence
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/μg)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μm filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_1107659](https://www.ebi.ac.uk/ols/ontologies/ab/term/AB_1107659)
Molecular Weight: 150 kDa

Description

The R6-5-D6 monoclonal antibody reacts with human CD54 also known as Intercellular Adhesion Molecule-1 (ICAM-1), is a single-chain transmembrane glycoprotein with a polypeptide core of 55 kDa and a member of the Ig superfamily. CD54 is typically expressed on non-hematopoietic cells such as endothelial cells, thymic epithelial cells, and fibroblasts but can also be expressed on macrophages, T-lymphoblasts, germinal center B cells and dendritic cells. CD54 is a ligand for LFA-1, a receptor found on leukocytes. When activated, leukocytes bind to endothelial cells via CD54/LFA-1 and then transmigrate into tissues. Pro-inflammatory cytokines such as IL-1, TNF and IFN γ induce CD54 expression on endothelial cells and fibroblasts. The R6-5-D6 antibody has been reported to inhibit the interaction of CD54 with leukocytes, thereby decreasing leukocyte adhesion to the vascular endothelium and inflammatory tissue injury.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0020-2#tab_references or scan the QR code below.



Bio X Cell, LLC
<https://bioxcell.com>
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC