

# Technical Data Sheet

## InVivoMAb anti-mouse L-Selectin (CD62L)



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** BE0021  
**Clone:** Mel-14  
**Isotype:** Rat IgG2a,  $\kappa$   
**Recommended Isotype Control(s):** InVivoMAb rat IgG2a isotype control, anti-trinitrophenol  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** C3H/eb mouse B lymphoma 38C-13  
**Reported Applications:** *in vivo* CD62L neutralization  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <2EU/mg (<0.002EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2  $\mu$ m filtered  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Purification:** Protein G  
**RRID:** [AB\\_1107665](https://abnova.com/AB_1107665)  
**Molecular Weight:** 150 kDa

### Description

The Mel-14 monoclonal antibody reacts with mouse CD62L also known as L-selectin and MEL-14. CD62L is a 76 kDa glycoprotein and a member of the selectin family that is expressed by neutrophils, monocytes, the majority of naïve T and B cells, a subset of memory T cells, NK cells, and most thymocytes. CD62L is a cell adhesion molecule that binds to many glycoprotein ligands including CD34, GlyCAM-1, MAdCAM-1, and PSGL-1 and acts as a "homing receptor" for lymphocytes to enter secondary lymphoid tissues via high endothelial venules. The Mel-14 monoclonal antibody reacts with mouse CD62L also known as L-selectin and MEL-14. CD62L is a 76 kDa glycoprotein and a member of the selectin family that is expressed by neutrophils, monocytes, the majority of naïve T and B cells, a subset of memory T cells, NK cells, and most thymocytes. CD62L is a cell adhesion molecule that binds to many glycoprotein ligands including CD34, GlyCAM-1, MAdCAM-1, and PSGL-1 and acts as a "homing receptor" for lymphocytes to enter secondary lymphoid tissues via high endothelial venules.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

## Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/catalogsearch/result/?q=BE0021#tab\\_references](https://bioxcell.com/catalogsearch/result/?q=BE0021#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**

