

InVivoMAb anti-mouse PD-1 (CD279)

Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0033-2
Clone:	J43
Isotype:	Armenian Hamster IgG
Recommended Isotype Control(s):	InVivoMAb polyclonal Armenian hamster IgG
Recommended Dilution Buffer:	InVivoPure pH 6.5 Dilution Buffer
Immunogen:	Syrian Hamster BKH cells transfected with mouse PD-1 cDNA <i>in vivo</i> blocking of PD-1/PD-L signaling <i>in vitro</i> PD-1 neutralization Western blot
Reported Applications:	
Formulation:	PBS, pH 6.5 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 μM filtered
Production:	Purified from tissue culture supernatant in an animal free facility
Purification:	Protein G
RRID:	AB_1107747
Molecular Weight:	150 kDa

Description

The J43 monoclonal antibody reacts with mouse PD-1 (programmed death-1) also known as CD279. PD-1 is a 50-55 kDa cell surface receptor encoded by the *Pdcd1* gene that belongs to the CD28 family of the immunoglobulin superfamily. PD-1 is transiently expressed on CD4 and CD8 thymocytes as well as activated T and B lymphocytes and myeloid cells. PD-1 expression declines after successful elimination of antigen. Additionally, *Pdcd1* mRNA is expressed in developing B lymphocytes during the pro-B-cell stage. PD-1's structure includes a ITIM (immunoreceptor tyrosine-based inhibitory motif) suggesting that PD-1 negatively regulates TCR signals. PD-1 signals via binding its two ligands, PD-L1 and PD-L2 both members of the B7 family. Upon ligand binding, PD-1 signaling inhibits T-cell activation, leading to reduced proliferation, cytokine production, and T-cell death. Additionally, PD-1 is known to play key roles in peripheral tolerance and prevention of autoimmune disease in mice as PD-1 knockout animals show dilated cardiomyopathy, splenomegaly, and loss of peripheral tolerance. Induced PD-L1 expression is common in many tumors including squamous cell carcinoma, colon adenocarcinoma, and breast adenocarcinoma. PD-L1 overexpression results in increased resistance of tumor cells to CD8 T cell mediated lysis. In mouse models of melanoma, tumor growth can be transiently arrested via treatment with antibodies which block the interaction between PD-L1 and its receptor PD-1. For these reasons anti-PD-1 mediated immunotherapies are currently being explored as cancer treatments. The J43 antibody has been shown to block the binding of both mouse PD-L1-Ig and mouse PD-L2-Ig to PD-1.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com


Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Binding Validation

Western blot data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, email technicalservice@bxccl.com.



Bio X Cell, Inc.

bxccl.com
1.866.787.3444

customerservice@bxccl.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell