Technical Data Sheet

InVivoMAb anti-mouse CD11c



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0038
Clone: N418

Isotype: Armenian Hamster lgG2

Recommended Isotype Control(s): InVivoMAb polyclonal Armenian hamster IqG

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer Immunogen: Mouse spleen dendritic cells In vivo targeting of dendritic cells

Functional assays Flow cytometry

Immunohistochemistry (frozen)
Immunohistochemistry (paraffin)

Immunofluorescence Immunoprecipitation

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein A

Molecular Weight: 150 kDa

Description

The N418 monoclonal antibody reacts with mouse CD11c, the most widely used defining marker for murine dendritic cells (DCs). CD11c is also known as integrin alpha-X (Itgax) and CD11 antigen-like family member C. CD11c is a 150-kDa single-pass type I membrane protein of the integrin alpha chain family, and its related members include CD11a (LFA-1), CD11b (MAC-1), and CD11d (D). In the context of sequence homology, expression profiles, and ligands, CD11c is substantially like CD11b and CD11d. CD11c is expressed on the plasma membranes of most DCs, monocytes, tissue macrophages, NK cells, and at a low level in neutrophils. CD11c binds with cell adhesion molecules (e.g., ICAM-1, ICAM-4), LPS from bacterial cell walls, iC3b complement protein, and fibrinogen, as well as collagen. CD11c's biological functions are not well studied, and pioneering studies showed it to mediate phagocytosis of iC3b-opsonized particles in vitro, which gave it recognition as complement receptor 4 (Cr4). CD11c is suggested to play a role in antigen presentation by the DCs and to mediate inflammatory responses in vivo. Due to its high expression on immature antigen DCs, CD11c is often considered an extremely effective immunotarget in experimental studies. The N418 antibody has also been demonstrated for targeting antigens to DCs (i.e., via CD11c), for rapid antibody generation from low immunogenic antigen targets, for anti-tumoral

Bio X Cell, LLC Page 1 of 2

immunomodulation, and for vaccine strategies. The N418 monoclonal antibody specifically binds on the surface of mouse DCs, and it does not bind peritoneal macrophages, B cells, or lymph node cells. The N418 antibody does not block dendritic cell binding to T cells.

Storage

Store at the stock concentration at 4°C . Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0038?bxcs=9k1b3a#tab_references or scan the QR code below



Bio X Cell, LLC https://bioxcell.com

+1-866-787-3444 customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2