# **Technical Data Sheet**

## InVivoMAb anti-mouse/human VLA-4 (CD49d)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

## Lot Specific Information

Lot Number: Lot Specific\* Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### **Product Information**

Catalog Number: BE0071 Clone: PS/2

**Isotype:** Rat IgG2b, κ

Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

Recommended Dilution Buffer: InVivoPure pH 6.5 Dilution Buffer

Immunogen:Mouse P815 mast cellsReported Applications:in vivo VLA-4 neutralizationin vitro VLA-4 neutralization

in vitro VLA-4 neutraliza Flow cytometry

**Formulation:** PBS, pH 6.5

Contains no stabilizers or preservatives

**Endotoxin:** <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

**Production:** Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB\_1107657

Molecular Weight: 150 kDa

#### **Description**

The PS/2 monoclonal antibody reacts with human and mouse VLA-4  $\alpha$  chain also known as CD49d and integrin alpha 4. VLA-4 is a 150 kDa glycoprotein belonging to the integrin family that is expressed by many cell types including T and B lymphocytes, monocytes, eosinophils, basophils, mast cells, thymocytes, NK cells, and dendritic cells. Integrin  $\alpha$ 4 associates with integrin  $\beta$ 7 to form integrin  $\alpha$ 4 $\beta$ 7 also known as LPAM-1 as well as integrin  $\beta$ 1 (CD29) to form integrin  $\alpha$ 4 $\beta$ 1 also known as VLA-4. Integrin  $\alpha$ 4 plays roles in adhesion and T cell co-stimulation. Integrin  $\alpha$ 4 ligands include VCAM-1, MAdCAM-1, and fibronectin. The PS/2 antibody is useful for in vivo and in vitro VLA-4 neutralization.

### **Storage**

Store at the stock concentration at 4°C. Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/fags">https://bioxcell.com/fags</a>.

#### **Protocol Information**

Bio X Cell, LLC Page 1 of 2

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### **Application References**

For a complete list of references, visit <a href="https://bioxcell.com/catalogsearch/result/?q=BE0071#tab\_references">https://bioxcell.com/catalogsearch/result/?q=BE0071#tab\_references</a> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2