

# Technical Data Sheet

## InVivoMAb anti-mouse TIM-2



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** BE0114  
**Clone:** RMT2-29  
**Isotype:** Rat Ig2a,  $\lambda$   
**Recommended Isotype Control(s):** InVivoMAb rat IgG2a isotype control, anti-trinitrophenol  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** Not available or unknown  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <2EU/mg (<0.002EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2  $\mu$ m filtered  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Purification:** Protein G  
**RRID:** [AB\\_10949464](https://eutils.ncbi.nlm.nih.gov/entrez/eutils/rrid.cgi?db=AB&term=AB_10949464)

### Description

The RMT2-29 monoclonal antibody reacts with mouse T cell immunoglobulin and mucin domain 2 (TIM-2), a type I cell-surface glycoprotein and member of the Ig superfamily. TIM-2 is preferentially expressed on TH2 cells as well as activated T cells. The TIM gene family, plays critical roles in regulating the immune response to viral infection. TIM-2 is also involved in the development of atopic disease and other TH2-biased immune responses.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### Application References

For a complete list of references, visit [https://bioxcell.com/catalogsearch/result/?q=BE0114#tab\\_references](https://bioxcell.com/catalogsearch/result/?q=BE0114#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**