

# Technical Data Sheet

## InVivoMAb anti-rat Kappa Immunoglobulin Light Chain



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

Lot Number: Lot Specific\*  
Volume: Lot Specific\*  
Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*  
Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

Catalog Number: **BE0122**  
Clone: **MAR 18.5**  
Isotype: Mouse IgG2a,  $\kappa$   
Recommended Isotype Control(s): InVivoMAb mouse IgG2a isotype control, unknown specificity  
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer  
Immunogen: Soluble rat immunoglobulin  
Reported Applications: *in vivo* B cell depletion in combination with anti-CD19 (clone [1D3](#)) and anti-CD22 (clone [Cy34.1](#))  
Formulation: PBS, pH 7.0  
Contains no stabilizers or preservatives  
Endotoxin: <2EU/mg (<0.002EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
Purity: >95%  
Determined by SDS-PAGE  
Sterility: 0.2  $\mu$ m filtered  
Production: Purified from cell culture supernatant in an animal-free facility  
Purification: Protein G  
RRID: [AB\\_10951292](#)  
Molecular Weight: 150 kDa

### Description

The MAR 18.5 monoclonal antibody reacts with the kappa chain of the rat immunoglobulin light chain. The  $\kappa$  chain is one of two types of polypeptide subunits which make up the immunoglobulin light chain. A typical antibody is composed of two immunoglobulin heavy chains and two immunoglobulin light chains. The  $\kappa$  chain is coded for by V (variable), J (joining) and C (constant) genes. These genes undergo V(D)J recombination to generate a diverse repertoire of immunoglobulins. This antibody is used in combination with rat anti-mouse CD19 and CD22 (clones [1D3](#) and [Cy34.1](#)) to deplete B cells *in vivo*.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/catalogsearch/result/?q=BE0122#tab\\_references](https://bioxcell.com/catalogsearch/result/?q=BE0122#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**