

Technical Data Sheet

InVivoMAb anti-mouse V γ 2 TCR



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0168
Clone: UC3-10A6
Isotype: Armenian Hamster IgG, κ
Recommended Isotype Control(s): InVivoMAb polyclonal Armenian hamster IgG
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: G8 mouse T cells
Reported Applications: *in vivo* $\gamma\delta$ T cell depletion
Flow cytometry
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_10950109](https://abnova.com/AB_10950109)
Molecular Weight: 150 kDa

Description

The UC3-10A6 monoclonal antibody reacts with an epitope on the delta chain of the mouse V γ 2 TCR (V gamma 2 T cell receptor). V γ 2 TCR expressing T lymphocytes make up a large proportion of the $\gamma\delta$ T cells in late fetal and adult thymus, peripheral lymphoid tissues, lung, intestinal epithelium, and epidermis. The exact function, ligand, and specificity of $\gamma\delta$ TCR-expressing T cells are not completely understood. Studies suggest that these cells recognize bacterial ligands and some tumor cells in the context of MHC class I-like gene products and play a role in regulating the immune response during bacterial infection. The UC3-10A6 antibody has been shown to deplete $\gamma\delta$ T cells when administered *in vivo*.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0168#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC