Technical Data Sheet

InVivoMAb anti-mouse Kappa Immunoglobulin Light Chain



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0176

 Clone:
 187.1 (HB-58)

 Isotype:
 Rat IgG1, κ

Recommended Isotype Control(s): InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: Mouse lgG2b Isotype control antibody clone MPC-11

Reported Applications: Immunofluorescence

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2ΕU/mg (<0.002ΕU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

Production: Purified from cell culture supernatant in an animal-free facility

 Purification:
 Protein G

 RRID:
 AB_10948999

 Molecular Weight:
 150 kDa

Description

The 187.1 monoclonal antibody reacts with the kappa chain of the mouse immunoglobulin light chain. The κ chain is one of two types of polypeptide subunits which make up the immunoglobulin light chain. A typical antibody is composed of two immunoglobulin heavy chains and two immunoglobulin light chains. The κ chain is coded for by V (variable), J (joining) and C (constant) genes. These genes undergo V(D)J recombination to generate a diverse repertoire of immunoglobulins.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Bio X Cell, LLC Page 1 of 2

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0176#tab_references or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2