

Technical Data Sheet

InVivoMAb anti-mouse IL-4



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0199
Clone: BVD6-24G2
Isotype: Rat IgG1, κ
Recommended Isotype Control(s): InVivoMAb rat IgG1 isotype control, anti-horseradish peroxidase
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Recombinant mouse IL-4
Reported Applications: ELISA
ELISPOT
Flow cytometry
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_10949184](https://rrid.info/AB_10949184)
Molecular Weight: 150 kDa

Description

The BVD6-24G2 monoclonal antibody reacts with mouse IL-4 (interleukin-4) a multifunctional 14 kDa cytokine. IL-4 is expressed primarily by activated Th2 cells and NK cells, and at lower levels by mast cells, and basophils. IL-4 signals through the IL-4R α . Upon receptor binding IL-4 stimulates activated B and T lymphocyte proliferation, and the differentiation of B cells into plasma cells. It also induces B cell class switching to IgE, and up-regulates MHC class II production while decreasing the production of Th1 cells, macrophages, IFN γ , and dendritic cell IL-12. Like other Th2 associated cytokines, IL-4 is involved in the airway inflammation observed in the lungs of patients with allergic asthma. The BVD6-24G2 antibody cannot neutralize the bioactivity of natural or recombinant IL-4.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result?q=BE0199#tab_references or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC
<https://bioxcell.com>
+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC