Technical Data Sheet

InVivoMAb anti-mouse F4/80



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <u>https://bioxcell.com/terms-and-conditions</u>.

Lot Specific Information

Lot Number:	Lot Specific*	
Volume:	Lot Specific*	
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *	
Total Protein:	Lot Specific*	
*This information will be noted on the certificate of analysis that ships with this product.		

Product Information

Catalog Number:	BE0206
Clone:	CI:A3-1
Isotype:	Rat lgG2b, к
Recommended Isotype Control(s):	InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	C57BL/6 mouse thioglycollate stimulated peritoneal macrophages
Reported Applications:	<i>in vivo</i> Monocyte/Macrophage depletion Functional assays Immunohistochemistry (paraffin) Immunohistochemistry (frozen) Flow cytometry
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/µg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 µm filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	<u>AB_10949019</u>
Molecular Weight:	150 kDa

Description

The CI:A3-1 monoclonal antibody reacts with mouse F4/80 also known as EMR1. F4/80 is a 160 kDa membrane glycoprotein belonging to the EGF-TM7 family. F4/80 is expressed on most tissue macrophages including peritoneal, lung, gut, thymus and splenic red pulp macrophages, as well as Kuffer cells, Langerhans cells, bone marrow stromal cells, and a subset of dendritic cells. F4/80's cognate ligand has not been identified however studies in mice have shown that F4/80 is important for cytokine production and induction regulatory T cells.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <u>https://bioxcell.com/catalogsearch/result/?q=BE0206#tab_references</u> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures. Not for resale. Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC