

InVivoMAb anti-mouse BTLA (CD272)



Lot Specific Information

| | |
|-----------------------|---|
| Lot Number: | Lot Specific* |
| Volume: | Lot Specific* |
| Concentration: | Lot Specific* (generally 4 to 11 mg/ml) * |
| Total Protein: | Lot Specific* |

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

| | |
|--|---|
| Catalog Number: | BE0210 |
| Clone: | 8F4 |
| Isotype: | Mouse IgG1, κ |
| Recommended Isotype Control(s): | InVivoMAb mouse IgG1 isotype control, unknown specificity |
| Recommended Dilution Buffer: | InVivoPure pH 7.0 Dilution Buffer |
| Immunogen: | C57BL/6 mouse BTLA Ig domain |
| Reported Applications: | Flow cytometry |
| Formulation: | PBS, pH 7.0 Contains no stabilizers or preservatives |
| Endotoxin: | <2EU/mg (<0.002EU/ μ g) Determined by LAL gel clotting assay |
| Purity: | >95% Determined by SDS-PAGE |
| Sterility: | 0.2 μ m filtered |
| Production: | Purified from tissue culture supernatant in an animal free facility |
| Purification: | Protein G |
| RRID: | AB_10948994 |
| Molecular Weight: | 150 kDa |

Description

The 8F4 monoclonal antibody reacts with mouse B- and T-lymphocyte attenuator (BTLA) also known as CD272. BTLA is an Ig superfamily member which is expressed on B cells, T cells, macrophages, dendritic cells, NK cells, and NKT cells. Like PD-1 and CTLA-4, BTLA interacts with a B7 homolog, B7-H4. However, unlike PD-1 and CTLA-4, BTLA displays T cell inhibition via interaction with tumor necrosis family receptors, not just the B7 family of cell surface receptors. BTLA is a ligand for herpes virus entry mediator (HVEM). BTLA-HVEM complexes have been shown to negatively regulate T cell immune responses. The 8F4 antibody reacts with BALb/c and C57BL/6 mouse BLTA.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**
All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <https://bxcell.com/product/m-cd272-btla/#references> or scan the QR code below.



Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell