

# Technical Data Sheet

## InVivoMAb anti-mouse Galectin-9



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** BE0218  
**Clone:** RG9-1  
**Isotype:** Rat IgG2b,  $\kappa$   
**Recommended Isotype Control(s):** InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** Recombinant mouse galectin-9  
**Reported Applications:** *in vivo* Galectin-9 blockade *in vitro* Galectin-9 blockade  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <2EU/mg (<0.002EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2  $\mu$ m filtered  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Purification:** Protein G  
**RRID:** [AB\\_2687702](https://abnova.com/AB_2687702)  
**Molecular Weight:** 150 kDa

### Description

The RG9-1 monoclonal antibody reacts with mouse galectin-9 a 40 kDa S-type lectin that is expressed by various cell types including lymphocytes, thymocytes, macrophages, dendritic cells, astrocytes, mast cells, eosinophils, fibroblasts, epithelial cells, and endothelial cells. Galectin-9 binds to  $\beta$ -galactosides and can serve as a ligand for TIM-3 (CD366). The protein is implicated in both innate and adaptive immune responses specifically, induction of cytokine secretion by macrophages, bactericidal functions, promotion of dendritic cell maturation, regulatory T cell expansion, and negative regulation of Th1, Th17, NK, and cytotoxic T cells. The RG9-1 antibody has been reported to block some biological activities of Galectin-9 when administered *in vivo*.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/catalogsearch/result/?q=BE0218#tab\\_references](https://bioxcell.com/catalogsearch/result/?q=BE0218#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**