Technical Data Sheet

InVivoMAb anti-human c-myc



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0238
Clone: 9E10
Isotype: Mouse IgG1

Recommended Isotype Control(s): InVivoMAb mouse IgG1 isotype control, unknown specificity

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: C-terminal peptide of human c-myc (aa 408-439)

Reported Applications: Western blot

ELISA

Immunoprecipitation Flow cytometry

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB_2687720

Molecular Weight: 150 kDa

Description

The 9E10 monoclonal antibody reacts with human c-myc, a 62 kDa transcription factor that plays a role in cell cycle progression, apoptosis and cellular transformation. Amplification of the c-myc gene has been found in several types of human cancers including lung, breast and colon carcinomas. c-Myc is commonly added to proteins of interest using recombinant DNA technology. The c-myc tag can then be used in many different assays that require recognition by an antibody.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Bio X Cell, LLC Page 1 of 2

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0238#tab_references or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2