

# Technical Data Sheet

## InVivoMAb anti-mouse 4-1BB (CD137)



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** BE0239  
**Clone:** 3H3  
**Isotype:** Rat IgG2a  
**Recommended Isotype Control(s):** InVivoMAb rat IgG2a isotype control, anti-trinitrophenol  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** Mouse CD137 human Fc fusion protein  
**Reported Applications:** *in vivo* 4-1BB stimulation  
*in vitro* 4-1BB stimulation  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <2EU/mg (<0.002EU/μg)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2 μm filtration  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Purification:** Protein G  
**RRID:** [AB\\_2687721](https://abnova.com/AB_2687721)  
**Molecular Weight:** 150 kDa

### Description

The 3H3 monoclonal antibody reacts with mouse 4-1BB, a TNF receptor superfamily member also known as CD137. 4-1BB is a 39 kDa transmembrane protein expressed by T lymphocytes, NK cells, dendritic cells, granulocytes, and mast cells. Upon binding its ligand 4-1BBL, 4-1BB provides costimulatory signals to both CD4 and CD8 T cells through the activation of NF-κB, c-Jun and p38 downstream pathways. The importance of the 4-1BB pathway has been underscored in a number of diseases, including cancer. Agonistic anti-4-1BB antibodies have been reported to induce T cell mediated antitumor immunity. The 3H3 antibody is an agonistic antibody that has been shown to stimulate 4-1BB signaling *in vivo*.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


## Application References

For a complete list of references, visit [https://bioxcell.com/catalogsearch/result?q=BE0239#tab\\_references](https://bioxcell.com/catalogsearch/result?q=BE0239#tab_references) or scan the QR code below.



## Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail [technicalservice@bioxcell.com](mailto:technicalservice@bioxcell.com).



---

### Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**