

Technical Data Sheet

InVivoMAb anti-human CD44



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0262
Clone: Hermes-1
Isotype: Rat IgG2a, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Human CD44
Reported Applications: *in vivo* CD44 blockade in xenografts
in vitro CD44 blockade
Western blot
Immunofluorescence
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_2687741](https://eutils.ncbi.nlm.nih.gov/entrez/eutils/rrid.cgi?db=AB_2687741)
Molecular Weight: 150 kDa

Description

The Hermes-1 monoclonal antibody reacts with human CD44 also known as Hermes, HCAM, and Pgp-1. CD44 is an 80-95 kDa glycoprotein that is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As an adhesion molecule, CD44 participates in a wide variety of cellular functions including lymphocyte activation, recirculation and homing, and hematopoiesis. CD44 is a receptor for hyaluronic acid and can also interact with other ligands, such as osteopontin, collagens, and matrix metalloproteinases (MMPs). Additionally, CD44 is involved in tumor metastasis and targeting of CD44 by antibodies has been shown to reduce the malignant activities of various neoplasms. Interestingly, high levels of the adhesion molecule CD44 on leukemic cells are essential to generate leukemia.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0262#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC