# **Technical Data Sheet**

InVivoMAb anti-mouse c-Kit (CD117)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

## Lot Specific Information

Lot Number: Lot Specific\* Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

## **Product Information**

Catalog Number: BE0280
Clone: 2B8

**Isotype:** Rat IgG2b, κ

Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer Immunogen: InvivoPure pH 7.0 Dilution Buffer

**Reported Applications:** Flow cytometry

Immunofluorescence Immunohistochemistry

**Formulation:** PBS, pH 7.0

Contains no stabilizers or preservatives

**Endotoxin:** <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

**Production:** Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID: AB\_2687803

Molecular Weight: 150 kDa

#### **Description**

The 2B8 monoclonal antibody reacts with mouse c-Kit also known as CD117, Steel factor receptor, stem cell factor receptor, and mast cell growth factor. c-Kit is a 145 kDa transmembrane tyrosine kinase and an immunoglobulin superfamily member. c-Kit is expressed on hematopoietic progenitor cells, mast cells, and acute myeloid leukemia (AML) cells. The interaction of the c-Kit receptor and its ligand stem cell factor (SCF), promotes the proliferation and differentiation of hematopoietic progenitor cells.

### **Storage**

Store at the stock concentration at  $4^{\circ}\text{C}$ . Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/faqs">https://bioxcell.com/faqs</a>.

### **Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

Bio X Cell, LLC Page 1 of 2

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## **Application References**

For a complete list of references, visit <a href="https://bioxcell.com/catalogsearch/result/?q=BE0280#tab\_references">https://bioxcell.com/catalogsearch/result/?q=BE0280#tab\_references</a> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2