

## InVivoMAb anti-mouse IFN $\gamma$ R $\alpha$ (CD119)

### Lot Specific Information

<b>Lot Number:</b>	Lot Specific*
<b>Volume:</b>	Lot Specific*
<b>Concentration:</b>	Lot Specific* (generally 4 to 11 mg/ml) *
<b>Total Protein:</b>	Lot Specific*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

<b>Catalog Number:</b>	<b>BE0287</b>
<b>Clone:</b>	<b>2E2</b>
<b>Isotype:</b>	Armenian Hamster IgG
<b>Recommended Isotype Control(s):</b>	InVivoMAb polyclonal Armenian hamster IgG
<b>Recommended Dilution Buffer:</b>	InVivoPure pH 7.0 Dilution Buffer
<b>Immunogen:</b>	Purified soluble recombinant mouse IFN $\gamma$ R $\alpha$ chain
<b>Reported Applications:</b>	Western blot Immunoprecipitation Flow cytometry
<b>Formulation:</b>	PBS, pH 7.0 Contains no stabilizers or preservatives
<b>Endotoxin:</b>	<2EU/mg (<0.002EU/ $\mu$ g) Determined by LAL gel clotting assay
<b>Purity:</b>	>95% Determined by SDS-PAGE
<b>Sterility:</b>	0.2 $\mu$ M filtered
<b>Production:</b>	Purified from tissue culture supernatant in an animal free facility
<b>Purification:</b>	Protein A
<b>RRID:</b>	AB_2687810
<b>Molecular Weight:</b>	150 kDa

### Description

The 2E2 monoclonal antibody reacts with the mouse IFN $\gamma$ R (interferon gamma receptor)  $\alpha$  chain also known as CD119 and IFN $\gamma$  receptor 1. CD119 heterodimerizes with IFN $\gamma$  receptor 2 (AF-1) to form the IFN $\gamma$ R, a class II cytokine receptor. The IFN $\gamma$ R is expressed ubiquitously on almost all cell types with the exception of mature erythrocytes. The 2E2 antibody is reported as a non-neutralizing antibody; it does not block the binding of IFN $\gamma$  to the receptor.

### Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at [bxcell.com/faqs](https://bxcell.com/faqs).

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

### Application References

For a complete list of references, visit <https://bxcell.com/product/invivomab-anti-mouse-ifnr-cd119/#references> or scan the QR code below.

### Bio X Cell, Inc.

bxcell.com  
1.866.787.3444  
[customerservice@bxcell.com](mailto:customerservice@bxcell.com)

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



---

**Bio X Cell, Inc.**

bxcell.com  
1.866.787.3444

**customerservice@bxcell.com**

*Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.*

**Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell**