

InVivoMAb anti-mouse c-Kit (CD117)

Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	BE0293
Clone:	ACK2
Isotype:	Rat IgG2b
Recommended Isotype Control(s):	InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	IL-3-dependent mast cells <i>in vivo</i> mast cell depletion <i>in vivo</i> c-Kit ⁺ cell depletion <i>in vitro</i> c-Kit neutralization Immunoprecipitation Flow cytometry
Reported Applications:	
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/μg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 μM filtered
Production:	Purified from tissue culture supernatant in an animal free facility
Purification:	Protein G
RRID:	
Molecular Weight:	

Description

The ACK2 monoclonal antibody reacts with mouse c-Kit also known as CD117, Steel factor receptor, stem cell factor receptor, and mast cell growth factor. c-Kit is a 145 kDa transmembrane tyrosine kinase and an immunoglobulin superfamily member. c-Kit is expressed on hematopoietic progenitor cells, mast cells, and acute myeloid leukemia (AML) cells. The interaction of the c-Kit receptor and its ligand stem cell factor (SCF), promotes the proliferation and differentiation of hematopoietic progenitor cells. The ACK2 antibody has been reported to deplete c-Kit⁺ cells, including mast cells, when administered *in vivo* and neutralize c-Kit⁺ signaling when used *in vitro*. This antibody is reported to cross-react with rat c-Kit.

Shelf-life and Storage

Store at the stock concentration at 4°C. **Do not freeze.**

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit <https://bxcell.com/product/invivomab-anti-mouse-c-kit-cd117-2/#references> or scan the QR code below.

Bio X Cell, Inc.

bxcell.com
1.866.787.3444
customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Bio X Cell, Inc.

bxcell.com
1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell