Technical Data Sheet

InVivoMAb anti-mouse E-selectin (CD62E)



Lot Specific Information

 Lot Number:
 Lot Specific*

 Volume:
 Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:BE0294Clone:9A9Isotype:Rat IgG2B

Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: COS-expressed, recombinant mouse E-selectin

Reported Applications: in vivo E-selectin blockade in vitro E-selectin blockade

Immunohistochemistry (frozen)

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2ΕU/mg (<0.002ΕU/μg)

Determined by LAL gel clotting assay

Purity:

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from tissue culture supernatant in an animal free facility

Purification:Protein GRRID:AB_2687816Molecular Weight:150 kDa

Description

The 9A9 monoclonal antibody reacts with mouse E-selectin also known as CD62E, endothelial-leukocyte adhesion molecule 1 (ELAM-1), and leukocyte-endothelial cell adhesion molecule 2 (LECAM2). E-selectin is a 115 kDa type I transmembrane protein and a member of the selectin family of adhesion molecules. E-selectin is expressed on cytokine-activated endothelial cells. Along with L-selectin and P-selectin, E-selectin mediates the initial interactions of leukocytes and platelets with endothelial cells. E-selectin is thought to play a role in inflammation, tumor metastasis, and angiogenesis. The 9A9 antibody has been shown to reduce the migration efficiency of Th1 cells and block E-selectin-mediated rolling.

Shelf-life and Storage

Store at the stock concentration at 4°C. Do not freeze.

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bxcell.com/product/invivomab-anti-mouse-e-selectin-cd62e/#references or scan the QR code below.

Bio X Cell, Inc.

bxcell.com

1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.



Bio X Cell, Inc.

bxcell.com 1.866.787.3444

customerser vice @bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.