Technical Data Sheet

InVivoMAb anti-mouse CD122 (IL-2Rβ)



Lot Specific Information

Lot Number:Lot Specific*Volume:Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Reported Applications:

 Catalog Number:
 BE0298

 Clone:
 TM-Beta 1

 Isotype:
 Rat IgG2b, κ

Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

Recommended Dilution Buffer: InVivoPure pH 8.0 Dilution Buffer

Immunogen: Mouse IL-2Rβ transfected cell line (TART-mβ cells)

in vivo NK cell depletion in vitro IL-2R blockade Functional assays Flow cytometry

Formulation: PBS, pH 8.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/µg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from tissue culture supernatant in an animal free facility

Purification:Protein GRRID:AB_2687820Molecular Weight:150 kDa

Description

The TM-beta 1 monoclonal antibody reacts with mouse CD122 also known as the IL-2 receptor beta chain. CD122 is a 70-75 kDa subunit of the IL-2 receptor and the IL-15 receptor. CD122 is expressed on NK cells and at lower levels by T lymphocytes, B lymphocytes, monocytes, and macrophages. The IL-2R has been shown to play roles in lymphocyte differentiation, activation, and proliferation. In complex with IL-2Ra, IL-2R binds IL-2 with relatively low affinity however, when CD122 combines with IL-2Ra and the common gamma chain (CD132) the complex binds IL-2 with high affinity. The TM-beta 1 antibody is reported to inhibit binding of IL-2 to the IL-2R. Additionally, this antibody has been shown to deplete CD122 expressing NK cells with administered in vivo.

Shelf-life and Storage

Store at the stock concentration at 4°C. Do not freeze.

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bxcell.com/product/invivomab-anti-mouse-cd122-il-2r/#references or scan the QR code below.

Bio X Cell, Inc.

bxcell.com 1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Bio X Cell, Bio X Cell Logo and all other trademarks are the property of Bio X Cell, Inc. © 2020 Bio X Cell



Bio X Cell, Inc.

bxcell.com 1.866.787.3444