

## InVivoMAb anti-human/monkey MHC class II (HLA-DR)

**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

Lot Number: Lot Specific\*  
Volume: Lot Specific\*  
Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*  
Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

Catalog Number: BE0306  
Clone: L243  
Isotype: Mouse IgG2a,  $\kappa$   
Recommended Isotype Control(s): InVivoMAb mouse IgG2a isotype control, unknown specificity  
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer  
Immunogen: Human lymphoblastoid B cell line RPMI 8866.9  
Reported Applications: *in vitro* blocking of MHC class II HLA-DR  
HLA class II binding assay  
*in vitro* MHC class II HLA-DR expressing cell negative selection  
Western blot  
Flow cytometry  
Formulation: PBS, pH 7.0  
Contains no stabilizers or preservatives  
Endotoxin:  $\leq 1$  EU/mg ( $\leq 0.001$  EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
Purity:  $\geq 95\%$   
Determined by SDS-PAGE  
Sterility: 0.2  $\mu$ m filtration  
Production: Purified from cell culture supernatant in an animal-free facility  
Purification: Protein G  
RRID: [AB\\_2736986](https://eutils.ncbi.nlm.nih.gov/entrez/eutils/rrid.cgi?db=AB_2736986)  
Molecular Weight: 150 kDa

### Description

The L243 monoclonal antibody reacts with the human and monkey MHC class II, HLA-DR. HLA-DR is a transmembrane glycoprotein composed of an  $\alpha$  chain (36 kDa) and a  $\beta$  chain (27 kDa). HLA-DR is expressed primarily on antigen presenting cells such as B cells, monocytes, macrophages, thymic epithelial cells and activated T cells. HLA-DR is critical for the presentation of peptides to CD4+ T cells.

### Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/be0306?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/be0306?bxcs=9k1b3a#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC**