# **Technical Data Sheet**

#### InVivoMAb anti-mouse CD69



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

# Lot Specific Information

Lot Number: Lot Specific\* Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### **Product Information**

Catalog Number: BE0330
Clone: CD69.2.2
Isotype: Mouse IgG1, κ

Recommended Isotype Control(s): InVivoMAb mouse IgG1 isotype control, unknown specificity

**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer Immunogen: CD69+ murine 300-19 pre-B cells

**Reported Applications:** in vivo down-regulation of CD69 expression

Functional assays

**Formulation:** PBS, pH 7.0

Contains no stabilizers or preservatives

**Endotoxin:** <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

**Production:** Purified from cell culture supernatant in an animal-free facility

 Purification:
 Protein A

 RRID:
 AB\_2894750

 Molecular Weight:
 150 kDa

### **Description**

The CD69.2.2 monoclonal antibody reacts with mouse CD69, an 85 kDa type II C-type lectin. CD69 expression is rapidly induced upon cell activation in all leukocytes. CD69 is thought to be a negative regulator of the immune response in part through modulating the production of TGF-β. Studies in CD69-deficient mice have revealed that CD69 plays crucial roles in the pathogenesis of various inflammatory diseases including arthritis, asthma, and colitis. When administered in vivo the CD69.2.2 antibody causes CD69 to be internalized but does not deplete CD69+ cells. Anti-CD69-treated mice resemble CD69-/- mice to a remarkable degree. The CD69.2.2 antibody has been shown to activate NK cells in vivo and in vitro and promote anti-tumor responses.

#### Storage

Store at the stock concentration at 4°C. **Do not freeze.** 

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/fags">https://bioxcell.com/fags</a>.

### **Protocol Information**

Bio X Cell, LLC Page 1 of 2

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## **Application References**

For a complete list of references, visit <a href="https://bioxcell.com/be0330?bxcs=9k1b3a#tab\_references">https://bioxcell.com/be0330?bxcs=9k1b3a#tab\_references</a> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2