

# Technical Data Sheet

## InVivoMAb anti-human CD220 (Insulin Receptor)



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** BE0338  
**Clone:** IR 83-22  
**Isotype:** Mouse IgG1  
**Recommended Isotype Control(s):** InVivoMAb mouse IgG1 isotype control, unknown specificity  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** Human IM-9 lymphocytes  
**Reported Applications:** Western blot  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <2EU/mg (<0.002EU/μg)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2 μm filtration  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Purification:** Protein G  
**RRID:** [AB\\_2894758](https://abnova.com/AB_2894758)  
**Molecular Weight:** 150 kDa

### Description

The IR 83-22 monoclonal antibody reacts with human CD220, also known as insulin receptor. CD220 is a type I transmembrane receptor tyrosine kinase which, upon binding to insulin, initiates several cellular effects, including glucose uptake, cell growth, differentiation, and apoptosis. These effects primarily occur in skeletal muscle, fat, and the liver. Neuronal CD220 signaling has been shown to have an important role in energy homeostasis, reproduction, and the development of neurodegenerative diseases. The IR 83-22 antibody has been shown to inhibit approximately 80% of insulin binding to IM-9 insulin receptor expressing cells.

### Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration

experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/be0338?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/be0338?bxcs=9k1b3a#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**