# **Technical Data Sheet**

## InVivoMAb anti-mouse NKG2A<sup>B6</sup>



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

# Lot Specific Information

Lot Number: Lot Specific\* Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

## **Product Information**

Catalog Number: BE0339
Clone: 16A11

**Isotype:** Mouse IgG2b, κ

Recommended Isotype Control(s): InVivoMAb mouse IgG2b isotype control, unknown specificity

**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer

Immunogen: C57BL/6 mouse CD94/NKG2A transfected CHO cells

**Reported Applications:** Flow cytometry **Formulation:** PBS, pH 7.0

Contains no stabilizers or preservatives

**Endotoxin:** <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

**Production:** Purified from cell culture supernatant in an animal-free facility

 Purification:
 Protein A

 RRID:
 AB\_2894759

 Molecular Weight:
 150 kDa

#### **Description**

The 16A11 monoclonal antibody reacts with NKG2A from C57BL/6 mice, the antibody does not react with BALB/c or 129 mouse strains. NKG2A, also known as CD159a is a type II transmembrane glycoprotein which belongs to the killer cell lectin-like receptor (KLR) family. NKG2A is expressed on NK cells, NKT cells, and activated CD8 T cells. NKG2A forms a disulfide-bonded heterodimer with CD94 that can bind to non-classical MHC class I antigen Qa-1 on target cells and inhibit NK cell activation.

## **Storage**

Store at the stock concentration at 4°C. **Do not freeze.** 

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/faqs">https://bioxcell.com/faqs</a>.

## **Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Bio X Cell, LLC Page 1 of 2

# **Application References**

For a complete list of references, visit <a href="https://bioxcell.com/be0339?bxcs=9k1b3a#tab\_references">https://bioxcell.com/be0339?bxcs=9k1b3a#tab\_references</a> or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2