

Technical Data Sheet

InVivoMAb anti-mouse/rat/bovine VLDL-R



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0345
Clone: IgG-6A6
Isotype: Mouse IgG1, λ
Recommended Isotype Control(s): InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: A peptide corresponding to the 10 C-terminal AA's (SVVSTDDDLA) of rat VLDL-R coupled to KLH.
Reported Applications: Western blot
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_2894764](https://ab2894764)
Molecular Weight: 150 kDa

Description

The IgG-6A6 monoclonal antibody reacts with the cytoplasmic tail of mouse, rat, and bovine very low-density lipoprotein receptor (VLDL-R). IgG-6A6 has also been reported to cross-react with human VLDL-R. VLDL-R is a member of the low-density lipoprotein (LDL) receptor family. The VLDL-R binds to triglyceride (TG)-rich lipoproteins but not LDL. VLDL-R is expressed in fatty acid-active tissues including heart, skeletal muscle, fat, and brain. Macrophages and dendritic cells also express VLDL-R. In contrast to the LDL receptor, VLDL-R binds apolipoprotein (apo) E2/2 VLDL particles as well as apoE3/3 VLDL. Various functions of the VLDL-R have been reported in lipoprotein metabolism, metabolic syndrome/atherosclerosis, cardiac fatty acid metabolism, neuronal migration, angiogenesis/tumor growth, and asthma.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0345?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC