

Technical Data Sheet

InVivoMAb anti-human CD71 (TfR1)



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0370
Clone: T56/14
Isotype: Mouse IgG1, κ
Recommended Isotype Control(s): InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Human hematopoietic cell line K562
Reported Applications: Flow cytometry
Immunoprecipitation
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Purification: Protein G
RRID: [AB_2927507](https://abnova.com/AB_2927507)
Molecular Weight: 150 kDa

Description

The T56/14 monoclonal antibody reacts with human CD71 also known as transferrin receptor or TfR1. CD71 is a type II homodimeric transmembrane glycoprotein which is expressed on the surface of proliferating cells, reticulocytes, and erythroid precursors. CD71 plays a role in the control of cellular proliferation and is required for iron import from transferrin into cells. CD71 is expressed on malignant cells at high levels and its expression correlates with cancer progression. The T56/14 antibody is a non-neutralizing antibody that does not inhibit the binding of transferrin.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0370#tab_references or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC