

Technical Data Sheet

InVivoMAb anti-mouse CD16.2 (FcγRIV)



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0378
Clone: 9E9
Isotype: Armenian hamster IgG
Recommended Isotype Control(s): InVivoMAb polyclonal Armenian hamster IgG
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Mouse FcγRIV extracellular domain-mouse IgG1 Fc fusion protein
Reported Applications: *in vivo* CD16.2 blockade
in vitro CD16.2 blockade
Flow cytometry
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/μg)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μm filtration
Purification: Protein A
RRID: [AB_2927515](https://abnova.com/AB_2927515)
Molecular Weight: 150 kDa

Description

The 9E9 monoclonal antibody reacts with mouse CD16.2, also known as FcγRIV (Fc receptor, IgG, low affinity IV). CD16.2 is a member of the immunoglobulin superfamily and is expressed on monocytes, macrophages, dendritic cells, and neutrophils. Fcγ receptors are essential for IgG-dependent effector functions *in vivo*. CD16.2 requires the common Fcγ chain for expression and signaling. CD16.2 binds to IgG2a and IgG2b with intermediate affinity. IgG2a- and IgG2b-dependent effector functions are severely impaired in CD16.2 deficient mice. CD16.2 has also been reported to be a low-affinity IgE receptor for all IgE allotypes and promotes IgE-induced lung inflammation. The 9E9 antibody has been shown to inhibit cellular CD16.2 function.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0378?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC