Technical Data Sheet

InVivoMAb anti-mouse TSLP



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <u>https://bioxcell.com/terms-and-conditions</u>.

Lot Specific Information

Lot Number:	Lot Specific*
Volume:	Lot Specific*
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein:	Lot Specific*
*This information w	ill be noted on the certificate of analysis that ships with this product.

Product Information

BE0379
28F12
Rat lgG2a, к
InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
InVivoPure pH 7.0 Dilution Buffer
Purified murine TSLP
<i>in vivo</i> TSLP neutralization <i>in vitro</i> TSLP neutralization
PBS, pH 7.0 Contains no stabilizers or preservatives
<2EU/mg (<0.002EU/µg) Determined by LAL gel clotting assay
>95% Determined by SDS-PAGE
0.2 µm filtration
Protein G
<u>AB_2927516</u>
150 kDa

Description

The 28F12 monoclonal antibody reacts with mouse Thymic Stromal-Derived Lymphopoetin (TSLP). TSLP is a cytokine that is closely related to IL-7. TSLP is expressed predominantly by epithelial cells in the thymus, lung, skin, intestine, and tonsils, as well as stromal cells, mast cells, and dendritic cells. TSLP exerts its biological activities by binding to a heterodimeric receptor that consists of IL-7R α and the TSLP receptor (TSLPR) chain. Expression of TSLP is regulated by NF- κ B and can be induced by exposure to viral, bacterial, or parasitic products, inflammatory cytokines, and the Th2 cell–associated cytokines IL-4 and IL-13. TSLP has important roles in conditioning DCs to drive Th2 differentiation and its expression is increased in immunopathologies associated with dysregulated Th2 cell–type cytokine production, including atopic dermatitis and asthma. It has additionally been reported to promote antibody isotype class switching in B cells, to regulate the development of Foxp3+ Tregs in the human thymus, and to promote basophil hematopoiesis. The 28F12 antibody has been reported to inhibit the bioactivity of mouse TSLP in vivo and in vitro.

Storage

Store at the stock concentration at 4°C. Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/? <u>q=BE0379#tab_references</u> or scan the QR code below.



Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail <u>technicalservice@bioxcell.com</u>.



Bio X Cell, LLC

https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC