

Technical Data Sheet

InVivoMAb anti-mouse VEGF-A



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: **BE0399**
Clone: **2G11-2A05**
Isotype: Rat IgG2a, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: N-terminal 24 AA sequence of murine VEGF
Reported Applications: *in vivo* VEGF-A neutralization
Western blot
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID:
Molecular Weight: 150 kDa

Description

The 2G11-2A05 monoclonal antibody reacts with mouse VEGF-A (vascular endothelial growth factor A). VEGF-A is a 45 kDa homodimeric, disulfide-linked glycoprotein that plays key roles in angiogenesis, vasculogenesis, and endothelial cell growth. It promotes mitogenesis, chemotaxis, and vascular permeability. Most tumors have a requirement for angiogenesis, and inhibition of VEGF binding to VEGF receptors has been the focus of several tumor therapeutic strategies. The 2G11-2A05 antibody neutralizes the biological activity of VEGF-A.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**


It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC