

# Technical Data Sheet

## InVivoMAb anti-human TROP-2



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

Lot Number: Lot Specific\*  
Volume: Lot Specific\*  
Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*  
Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

Catalog Number:	BE0408
Clone:	Pr1E11
Isotype:	Mouse IgG1, $\kappa$
Recommended Isotype Control(s):	InVivoMAb mouse IgG1 isotype control, unknown specificity
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Primary human prostate cancer cells
Reported Applications:	Western blot Immunohistochemistry (frozen) Flow cytometry
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<2EU/mg (<0.002EU/ $\mu$ g) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 $\mu$ m filtered
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	
Molecular Weight:	150 kDa

### Description

The Pr1E11 monoclonal antibody reacts with human TROP-2, also known as TACSTD2, EGP-1, and GA733-1. TROP-2 is a type I transmembrane glycoprotein with high homology to TROP-1/EpCAM. TROP-2 spans the epithelial membrane surface and plays a role in embryonic development, cell self-renewal, proliferation, and transformation. TROP-2 is found on the surface of multiple normal epithelial tissues, including skin and oral mucosa. TROP-2 can promote tumor growth and its overexpression is common in many types of malignant epithelial tumors. A variety of human epithelial cancer cells are characterized by TROP-2 overexpression, including breast, lung, urothelial, gastric, colorectal, pancreatic, prostatic, cervical, head and neck, and ovarian carcinomas.

### Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

### Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.


**Application References**

For a complete list of references, visit [https://bioxcell.com/be0408?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/be0408?bxcs=9k1b3a#tab_references) or scan the QR code below.



**Binding Validation**

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail [technicalservice@bioxcell.com](mailto:technicalservice@bioxcell.com).



**Bio X Cell, LLC**  
<https://bioxcell.com>  
+1-866-787-3444  
[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*  
*Not for resale.*  
**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC**