Technical Data Sheet

InVivoMAb anti-rat TCR α/β



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <u>https://bioxcell.com/terms-and-conditions</u>.

Lot Specific Information

| Lot Number: | Lot Specific* | |
|--|---|--|
| Volume: | Lot Specific* | |
| Concentration: | Lot Specific* (generally 4 to 11 mg/ml) * | |
| Total Protein: | Lot Specific* | |
| *This information will be noted on the certificate of analysis that ships with this product. | | |

Product Information

| Catalog Number: | BE0413 |
|---------------------------------|---|
| Clone: | R73 |
| Isotype: | Mouse lgG1, к |
| Recommended Isotype Control(s): | InVivoMAb mouse IgG1 isotype control, unknown specificity |
| Recommended Dilution Buffer: | InVivoPure pH 7.0 Dilution Buffer |
| Immunogen: | Rat T Blasts and Erythrocytes |
| Reported Applications: | in vivo administration |
| Formulation: | PBS, pH 7.0 Contains no stabilizers or preservatives |
| Endotoxin: | <2EU/mg (<0.002EU/µg) Determined by LAL gel clotting assay |
| Purity: | >95% Determined by SDS-PAGE |
| Sterility: | 0.2 µm filtration |
| Production: | Purified from cell culture supernatant in an animal-free facility |
| Purification: | Protein G |
| RRID: | |
| Molecular Weight: | 150 kDa |

Description

The R73 monoclonal antibody reacts with a framework component of rat α/β TCR (alpha beta T cell receptor). The α/β TCR is expressed by most peripheral T cells, intestinal epithelial lymphocytes, and thymocytes. The R73 antibody has been reported to suppress the development of experimental autoimmune encephalitis and adjuvant arthritis. Moreover, R73 treatment was applied in the field of transplantation and prevented allograft rejection in heart, skin, small bowel, kidney, and cornea.

Storage

Store at the stock concentration at 4°C. Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Bio X Cell, LLC

https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures. Not for resale. Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC