Technical Data Sheet

InVivoMAb anti-human VISTA



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific* Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0436 Clone: GG8

Isotype: Mouse IgG1, κ

Recommended Isotype Control(s): InVivoMAb mouse IgG1 isotype control, unknown specificity

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: Not available or unknown

Reported Applications: Immunohistochemistry-paraffin

Flow cytometry

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/µg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID:

Molecular Weight: 150 kDa

Description

The GG8 monoclonal antibody reacts with human VISTA (V-type immunoglobulin domain-containing suppressor of T cell activation), which is a negative checkpoint regulator (NCR) from the B7 family. Several in vitro and in vivo studies have demonstrated that the VISTA protein acts as a ligand as well as a receptor. VISTA is mainly expressed by myeloid cells and T cells, and in the CNS, VISTA is also expressed by microglial and endothelial cells. Unlike other NCR proteins, VISTA is expressed on naïve T cells, wherein its expression is negatively regulated by T cell activation. Besides its role in regulation of T cell quiescence (i.e., the suppression of T cells), VISTA is involved in efferocytosis, cytokine production (IL-10, IFN-gamma, and TNF-alpha), and chemotaxis in myeloid cells.

Storage

Store at the stock concentration at 4°C . Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate

Bio X Cell, LLC Page 1 of 2

dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0436?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2