

Technical Data Sheet

InVivoMAb anti-rat CD4



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0439
Clone: W3/25
Isotype: Mouse IgG1, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Reported Applications: *in vivo* down-regulation of surface CD4
in vitro neutralization of CD4
Flow cytometry
Immunohistochemistry (paraffin)
Immunohistochemistry (frozen)
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/ μ g)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μ m filtration
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID:
Molecular Weight: 150 kDa

Description

The W3/25 monoclonal antibody reacts with rat CD4. CD4 is a 55 kDa type I membrane glycoprotein from the immunoglobulin superfamily, and as a cell surface receptor protein, it acts as a co-receptor (with TCR) for class II MHC molecules displayed by antigen-presenting cells (APC). CD4 is expressed by the majority of thymocytes, helper T cells, a subset of NK-T cells, and weakly by dendritic cells and macrophages. CD4 plays an important role in the development of T cells and is required for the optimal functioning of mature T cells. The clone W3/25 is used as a "helper T-cell marker." The clone W3/25 is a nondepleting monoclonal antibody, but it has been shown to neutralize CD4 *in vitro*. Several *in vitro* studies also showed that the W3/25 monoclonal antibody inhibits CD4+ T cell activation by downregulating the CD4 molecule on the surface of lymphocytes. The W3/25 monoclonal antibody also inhibits antigen-induced T cell proliferation and IL2 production in MLR experiments. In prophylactic and experimental therapeutic *in vivo* investigations, this antibody has been extensively used to modulate disease-related and mechanistic parameters in experimental allergic encephalomyelitis (EAE), adjuvant arthritis, etc.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/catalogsearch/result/?q=BE0439#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC