

Technical Data Sheet

InVivoMAb anti-mouse CXCR4 (CD184)



bioxcell.com

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Website Link: <https://bioxcell.com/invivomab-anti-mouse-cxcr4-cd184-be0455>

Product Information

Catalog Number: BE0455
Clone: Cx4Mab-1
Isotype: Rat IgG2a, κ
Recommended Isotype Control(s): InVivoMAb rat IgG2a isotype control, anti-trinitrophenol
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: mCXCR4-PA tag overexpressing LN229 cells
Reported Applications: Flow cytometry
For details on *in vivo* applications please contact technicalservice@bioxcell.com
Formulation: PBS, pH 7.0
Contains no stabilizers or preservatives
Endotoxin: ≤ 1 EU/mg (≤ 0.001 EU/ μ g)
Determined by LAL assay
Purity: $\geq 95\%$
Determined by SDS-PAGE
Sterility: 0.2 μ m filtered
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID:
Molecular Weight: 150 kDa

Description

The Cx4Mab-1 monoclonal antibody reacts with mouse CXC chemokine receptor 4 (CXCR4), also called fusin and CD184. CXCR4 is a hepta-transmembrane domain G protein-coupled receptor (GPCR) that is expressed in hematopoietic cells, endothelial cells, neurons, and embryonic as well as adult stem cells. The primary ligand of CXCR4 is CXCL12, also known as stromal cell-derived factor-1 α (SDF-1 α). The CXCR4-CXCL12 signaling pathway plays a critical role in activating multiple signaling pathways including ERK1/2, ras, p38 MAPK, PLC/MAPK, and SAPK/JNK. CXCR4-CXCL12 signaling also regulates several biological processes including cell survival, proliferation, migration, and stemness. Overexpression of CXCR4 is often linked to poor prognosis in various cancers, and during cancer metastasis, CXCR4 positive cancer cells are chemotactically homed to tissues such as the liver, bone marrow, lungs, and lymph nodes that express abundant levels of CXCL12. CXCR4 as well as CXCR4/CXCL12 antagonists are emerging as attractive targets for experimental cancer therapeutics. The anti-mouse CXCR4 antibody, clone Cx4Mab-1, was generated using a cell-based immunization and screening approach. Its high affinity for both endogenous and exogenous mouse CXCR4 suggests its potential utility in *in vivo* preclinical mouse models.

Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out

of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/invivomab-anti-mouse-cxcr4-cd184-be0455?utm_source=cr9k1b#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © Bio X Cell, LLC