Technical Data Sheet

InVivoMAb anti-mouse ACKR4 (CCR11)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific* Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BE0467
Clone: A4Mab-3
Isotype: Rat IgG2b, κ

Recommended Isotype Control(s): InVivoMAb rat IgG2b isotype control, anti-keyhole limpet hemocyanin

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: A synthetic peptide corresponding to the N-terminal extracellular region of

mACKR4 (amino acids 1-19)

Reported Applications: Flow cytometry

ELISA

For details on in vivo applications, please contact

technicalservice@bioxcell.com

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <2EU/mg (<0.002EU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtered

Production: Purified from cell culture supernatant in an animal-free facility

Purification: Protein G

RRID:

Molecular Weight: 150 kDa

Description

The A4Mab-3 monoclonal antibody reacts with mouse atypical chemokine receptor 4 (ACKR4), also known as CCR11, CCRL1, CCX CKR, PPR1, and CCBP2. ACKR4 is expressed on T cells, stromal cells, and a subset of lymphatic endothelial cells within the skin, spleen, and gut. ACKR4 is a seven-transmembrane domain-containing protein that belongs to the atypical chemokine receptors (ACKRs) family. Due to their function, ACKRs are considered chemokine decoy receptors, internalizing receptors (interceptors), or chemokine-scavenging receptors. They mediate β-arrestin-dependent internalization of chemokines, followed by lysosomal degradation of the receptor-ligand complex. ACKR4 plays a key role in regulating cell migration by reducing the availability of inflammatory chemokines, specifically CCL19, CCL22, and CCL25, thereby limiting migratory responses mediated by CCR7, CCR6, CCR4, and CCR9. ACKR4 negatively regulates CXCR3-induced chemotaxis and facilitates the homing of CC7+ dendritic cells through scavenging and shaping CCL19 and CCL21 gradients in lymph nodes. ACKR4 is also involved in the regulation of thymic T-cell development and suppression of spontaneous autoimmunity. Loss of ACKR4 in colorectal cancer in mice is reported to impair DC migration to tumor-draining lymph nodes, which leads to a reduced number of tumor-specific T cells and resistance to immune checkpoint blockade therapy.

Bio X Cell, LLC Page 1 of 2

Storage

Store at the stock concentration at 4°C . Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/be0467?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC https://bioxcell.com

+1-866-787-3444
customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2025 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2