Technical Data Sheet

InVivoPlus anti-mouse IL-4



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bxcell.com/terms-and-conditions/.

Lot Specific Information

Lot Number: Lot Specific* Lot Specific* Volume:

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BP0045 Clone: 11R11 Rat IgG1, ĸ Isotype:

Recommended Isotype Control(s): InVivoPlus rat IgG1 isotype control, anti-horseradish peroxidase

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Immunogen: Partially purified native mouse IL-4

> in vivo IL-4 neutralization in vitro IL-4 neutralization

Reported Applications: in vivo IL-4 receptor stimulation (as a complex with IL-4)

Flow cytometry Western blot

PBS, pH 7.0 Formulation: Contains no stabilizers or preservatives

<1EU/mg (<0.001EU/µg)

Determined by LAL gel clotting assay

Purity:

Determined by SDS-PAGE

Sterility: 0.2 uM filtered

Production: Purified from tissue culture supernatant in an animal free facility

Purification: Protein G <5% **Aggregation:**

Determined by DLS RRID: AB_1107707 **Molecular Weight:** 150 kDa

Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative,

Description

Endotoxin:

The 11B11 monoclonal antibody reacts with mouse IL-4 (interleukin-4) a multifunctional 14 kDa cytokine. IL-4 is expressed primarily by activated Th2 cells and NK cells, and at lower levels by mast cells, and basophils. IL-4 signals through the IL-4Ra. Upon receptor binding IL-4 stimulates activated B and T lymphocyte proliferation, and the differentiation of B cells into plasma cells. It also induces B cell class switching to IgE, and up-regulates MHC class II production while decreasing the production of Th1 cells, macrophages, IFNy, and dendritic cell IL-12. Like other Th2 associated cytokines, IL-4 is involved in the airway inflammation observed in the lungs of patients with allergic asthma. The 11B11 monoclonal antibody has been shown to neutralize the bioactivity of natural or recombinant IL-4.

Shelf-life and Storage

Store at the stock concentration at 4°C. Do not freeze.

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on

Bio X Cell, Inc.

bxcell.com 1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

how to remove floccules or precipitates see our FAQ's at bxcell.com/faqs.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bxcell.com/product/invivoplus-anti-m-il-4/#references or scan the QR code below.



Bio X Cell, Inc.

bxcell.com

1.866.787.3444

customerservice@bxcell.com