Technical Data Sheet

InVivoPlus anti-mouse CD8a



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bxcell.com/terms-and-conditions/.

Lot Specific Information

Lot Number: Lot Specific* Lot Specific* Volume:

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: BP0117 YTS 169.4 Clone: Rat IgG2b, ĸ Isotype:

Recommended Isotype Control(s): InVivoPlus rat IgG2b isotype control, anti-keyhole limpet hemocyanin

InVivoPure pH 7.0 Dilution Buffer **Recommended Dilution Buffer:**

Immunogen: CBA mouse thymocytes

in vivo CD8+ T cell depletion **Reported Applications:**

Western blot PBS, pH 7.0

Formulation: Contains no stabilizers or preservatives

<1EU/mg (<0.001EU/µg)

Endotoxin: Determined by LAL gel clotting assay

>95%

Purity: Determined by SDS-PAGE

0.2 µM filtered Sterility:

Production: Purified from tissue culture supernatant in an animal free facility

Purification: Protein G <5%

Aggregation: Determined by DLS RRID: AB_10950145

Molecular Weight: 150 kDa

Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative,

Description

The YTS 169.4 monoclonal antibody reacts with mouse CD8a. The CD8 antigen is a transmembrane glycoprotein that acts as a co-receptor for the T cell receptor (TCR). Like the TCR, CD8 binds to class I MHC molecules displayed by antigen presenting cells (APC). CD8 is primarily expressed on the surface of cytotoxic T cells, but can also be found on thymocytes, natural killer cells, and some dendritic cell subsets. CD8 most commonly exists as a heterodimer composed of one CD8a and one CD8\beta chain however, it can also exist as a homodimer composed of two CD8\alpha chains. Both the CD8\alpha and CD8\beta chains share significant homology to immunoglobulin variable light chains. The molecular weight of each CD8 chain is approximately 34 kDa. The YTS 169.4 antibody exhibits depleting activity when used in vivo.

Shelf-life and Storage

Store at the stock concentration at 4°C. Do not freeze.

All Bio X Cell antibodies have a guaranteed shelf-life of one year from the date of customer receipt when stored as recommended. It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at bxcell.com/fags.

Bio X Cell, Inc.

bxcell.com 1.866.787.3444

customerservice@bxcell.com

Conditions: For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bxcell.com/product/invivoplus-anti-m-cd8/#references or scan the QR code below.



Bio X Cell, Inc.