Technical Data Sheet

InVivoPlus anti-mouse CTLA-4 (CD152)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <u>https://bioxcell.com/terms-and-conditions</u>.

Lot Specific Information

Lot Number:	Lot Specific*	
Volume:	Lot Specific*	
Concentration:	Lot Specific* (generally 4 to 11 mg/ml) *	
Total Protein:	Lot Specific*	
*This information will be noted on the certificate of analysis that ships with this product.		

Product Information

Catalog Number:	BP0131
Clone:	9H10
Isotype:	Syrian hamster IgG
Recommended Isotype Control(s):	InVivoPlus polyclonal Syrian hamster IgG
Recommended Dilution Buffer:	InVivoPure pH 7.0 Dilution Buffer
Immunogen:	Mouse CTLA-4-human IgG1 fusion protein
Reported Applications:	<i>in vivo</i> CTLA-4 neutralization <i>in vitro</i> CTLA-4 neutralization Western blot
Formulation:	PBS, pH 7.0 Contains no stabilizers or preservatives
Endotoxin:	<1EU/mg (<0.001EU/µg) Determined by LAL gel clotting assay
Purity:	>95% Determined by SDS-PAGE
Sterility:	0.2 µm filtration
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
Aggregation:	<5% Determined by SEC
RRID:	<u>AB_10950184</u>
Molecular Weight:	150 kDa

Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

Description

The 9H10 monoclonal antibody reacts with mouse CTLA-4 (cytotoxic T lymphocyte antigen-4) also known as CD152. CTLA-4 is a 33 kDa cell surface receptor encoded by the Ctla4 gene that belongs to the CD28 family of the lg superfamily. CTLA-4 is expressed on activated T and B lymphocytes. CTLA-4 is structurally similar to the T-cell co-stimulatory protein, CD28, and both molecules bind to the B7 family members B7-1 (CD80) and B7-2 (CD86). Upon ligand binding, CTLA-4 negatively regulates cell-mediated immune responses. CTLA-4 plays roles in induction and/or maintenance of immunological tolerance,

thymocyte development, and regulation of protective immunity. The critical role of CTLA-4 in immune down-regulation has been demonstrated in CTLA-4 deficient mice, which succumb at 3-5 weeks of age due to the development of a lymphoproliferative disease. CTLA-4 is among a group of inhibitory receptors being explored as cancer treatment targets through immune checkpoint blockade. The 9H10 antibody has been shown to promote T cell co-stimulation by blocking CTLA-4 binding to the B7 co-receptors, allowing for CD28 binding.

Storage

Store at the stock concentration at 4°C. Do not freeze.

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/faqs.

Protocol Information


Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Binding Validation

Application References

For a complete list of references, visit <u>https://bioxcell.com/bp0131?</u> <u>bxcs=9k1b3a#tab_references</u> or scan the QR code below.





antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com. <u>kDa</u> 1 2 3

Validation data shown below confirms that this clone binds to its target

Lane 1: 0.2 μ g reduced purified mouse CTLA-4 with histidine tag at C-terminus Lane 2: 0.1 μ g reduced purified mouse CTLA-4 with histidine tag at C-terminus Lane 3: 0.05 μ g reduced purified mouse CTLA-4 with histidine tag at C-terminus

Primary: biotinylated anti-mouse CTLA-4 (CD152) antibody (9H10) at 8 µg/ml

Secondary: HRP labeled streptavidin 1:1000 dilution

Predicted band size : 25-30 kDa

Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

, Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC @ 2024 Bio X Cell, LLC