# **Technical Data Sheet**

RecombiMAb anti-mouse CD40L (CD154)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <a href="https://bioxcell.com/terms-and-conditions">https://bioxcell.com/terms-and-conditions</a>.

## **Lot Specific Information**

Lot Number: Lot Specific\* Volume: Lot Specific\*

Concentration: Lot Specific\* (generally 4 to 11 mg/ml) \*

Total Protein: Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

#### **Product Information**

Catalog Number: CP033

Clone: MR-1-CP033 Isotype: Mouse IgG2a, κ

Recommended Isotype Control(s): InVivoPlus mouse IgG2a isotype control, unknown specificity

**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer Immunogen: Activated mouse Th1 clone D1.6

**Reported Applications:** in vivo blocking of CD40/CD40L signaling\*

in vitro blocking of CD40/CD40L signaling\*

Western blot\*

\*Reported for the original Armenian hamster IgG MR-1 antibody

**Formulation:** PBS, pH 7.0

Contains no stabilizers or preservatives

**Endotoxin:** <1EU/mg (<0.001EU/µg)

Determined by LAL gel clotting assay

**Purity:** >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

**Production:** Purified from HEK293 cell supernatant in an animal-free facility

Purification: Protein G
Aggregation: <5%

Determined by SEC

**RRID:** 

Molecular Weight: 150 kDa

### **Murine Pathogen Test Results**

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

### **Description**

The MR-1-CP033 monoclonal antibody is a chimeric version of the original MR-1 antibody. The variable domain sequences are identical to the original MR-1 but the constant region sequences have been switched from Armenian hamster IgG to mouse IgG2a. The MR-1-CP033 antibody contains no Fc mutations just as the original Armenian hamster IgG antibody does not. The MR-1-CP033 antibody reacts with mouse CD154 also known as CD40 ligand. CD154 exists as a 39 kDa

Bio X Cell, LLC Page 1 of 2

accessory molecule and belongs to the TNF superfamily of cytokines. CD154 is primarily expressed on the surface of activated CD4+ T lymphocytes but can also be expressed by platelets, mast cells, macrophages, basophils, NK cells, B lymphocytes, CD8+ T lymphocytes as well as non-hematopoietic cells including smooth muscle cells, endothelial cells, and epithelial cells. CD154 signals through CD40 and is thought to play a key role in T and B lymphocyte costimulation. The MR-1 monoclonal antibody has been reported to inhibit in vitro activation of B lymphocytes by blocking the binding of CD154 with CD40 on T helper cells as well as inhibit the formation of germinal centers and disrupt antigen-specific T cell responses. Additionally, the MR-1 antibody blocks interactions of T cells and antigen-presenting cells in vitro and blocks the development of experimental autoimmune disease in vivo.

## **Storage**

Store at the stock concentration at 4°C. **Do not freeze.** 

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <a href="https://bioxcell.com/fags">https://bioxcell.com/fags</a>.

#### **Protocol Information**

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Bio X Cell, LLC

https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2