Technical Data Sheet

RecombiMAb anti-mouse ICOSL (CD275) (D265A)



<u>Attention</u>: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number: Lot Specific* Volume: Lot Specific*

Concentration: Lot Specific* (generally 4 to 11 mg/ml) *

Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: CP044

Clone: HK5.3-CP044
Isotype: Mouse IgG1, κ

Recommended Isotype Control(s): RecombiMAb mouse IgG1 (D265A) isotype control, anti-hen egg lysozyme

Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer

Mutations: D265A

Immunogen: Mouse B7-RP1 transfected cell line

Reported Applications: in vivo ICOSL neutralization*

*Reported for the original rat lgG2a HK5.3 antibody

Formulation: PBS, pH 7.0

Contains no stabilizers or preservatives

Endotoxin: <1EU/mg (<0.001EU/μg)

Determined by LAL gel clotting assay

Purity: >95%

Determined by SDS-PAGE

Sterility: 0.2 µm filtration

Production: Purified from HEK293 cell supernatant in an animal-free facility

Purification: Protein G
Aggregation: <5%

Determined by SEC

RRID:

Molecular Weight: 150 kDa

Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

Description

The HK5.3-CP044 monoclonal antibody is a chimeric version of the original HK5.3 antibody. The variable domain sequences are identical to the original HK5.3 but the constant region sequences have been switched from rat lgG2a to mouse lgG1. The HK5.3-CP044 antibody also contains a D265A mutation in the Fc fragment rendering it unable to bind to endogenous Fcγ receptors. The HK5.3-CP044 monoclonal antibody reacts with mouse ICOSL (inducible T cell co-stimulator ligand) also known as CD275, B7RP-1, and B7-H2. ICOSL is a 40 kDa immune checkpoint protein belonging to the lg

Bio X Cell, LLC Page 1 of 2

receptor superfamily. Upon ICOSL binding, ICOS signaling co-stimulates T and B cell responses. The ligand Is expressed on antigen presenting cells including splenic B cells, dendritic cells, and macrophages. ICOS signaling is also thought to be important for maintaining regulatory T cell homeostasis. The HK5.3 antibody has been shown to block the binding of ICOSL to ICOS both in vitro and in vivo. HK5.3 treatment of mice has been reported to lead to a loss of regulatory T cells.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at https://bioxcell.com/fags.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Bio X Cell, LLC https://bioxcell.com +1-866-787-3444 customerservice@bioxcell.com Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC

Bio X Cell, LLC Page 2 of 2