

# Technical Data Sheet

## RecombiMAb anti-mouse CD3ε



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** CP082  
**Clone:** 145-2C11-CP082  
**Isotype:** Mouse IgG2a, κ  
**Recommended Isotype Control(s):** RecombiMAb mouse IgG2a isotype control, anti-hen egg lysozyme  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Immunogen:** Mouse BM10-37 cytotoxic T cells  
**Reported Applications:** *in vivo* T cell depletion  
*in vitro* T cell stimulation/activation  
*in vivo* T cell stimulation/activation  
Immunofluorescence  
Flow cytometry  
Western blot  
\*Reported for the original 145-2C11 antibody. For information on *in vivo* applications, please contact [technicalservice@bioxcell.com](mailto:technicalservice@bioxcell.com)

**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives

**Endotoxin:** ≤0.5EU/mg (≤0.0005EU/μg)  
Determined by LAL assay

**Purity:** ≥95%  
Determined by SDS-PAGE

**Sterility:** 0.2 μm filtration

**Production:** Purified from mammalian cell supernatant in an animal-free facility

**Purification:** Protein G

**Aggregation:** <5%  
Determined by SEC

**RRID:**  
**Molecular Weight:** 150 kDa

### Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

### Description

The 145-2C11-CP082 monoclonal antibody is a recombinant, Fc-engineered chimeric version of the original 145-2C11 antibody. The variable domain sequences are identical but the constant region sequences have been switched from hamster

IgG1 to mouse IgG2a, κ for use in murine models. Studies comparing the chimeric mouse IgG2a antibody to the original hamster isotype show both bind murine Fcγ receptors avidly and had similar activating, immunogenic, and immunosuppressive properties in mice. Species-matched chimeric antibodies demonstrate reduced immunogenicity and formation of anti-drug antibodies (ADAs) compared to xenogenic antibodies in animal models. The highly controlled sequence and lack of genetic drift in recombinant antibodies provide more reliable and reproducible results over hybridoma derived antibodies. The 145-2C11-CP082 monoclonal antibody reacts with mouse CD3ε, a 20 kDa transmembrane cell-surface protein that belongs to the immunoglobulin superfamily. CD3ε is one of five polypeptide chains that combine to form the TCR complex. CD3ε is expressed on T lymphocytes, NK-T cells, and to varying degrees on developing thymocytes. CD3 plays roles in TCR signaling, T lymphocyte activation, and antigen recognition. The 145-2C11 antibody has been shown to induce T lymphocyte activation, proliferation, and apoptosis via binding and stimulating the TCR. Additionally, the 145-2C11 antibody has been reported to block the binding of the 17A2 antibody to CD3ε+ T lymphocytes. References: Alegre, M L et al. "An anti-murine CD3 monoclonal antibody with a low affinity for Fc gamma receptors suppresses transplantation responses while minimizing acute toxicity and immunogenicity." Journal of immunology (Baltimore, Md. : 1950) vol. 155,3 (1995): 1544-55.

## Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

## Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/cp082?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/cp082?bxcs=9k1b3a#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**  
<https://bioxcell.com>  
+1-866-787-3444  
[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2026 Bio X Cell, LLC**