Technical Data Sheet

FlowMAb FITC anti-mouse Ly6G/Ly6C (Gr-1)

Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at https://bioxcell.com/terms-and-conditions.

Lot Specific Information

Lot Number:	**Lot Specific***
Volume:	**Lot Specific***
Concentration:	**Lot Specific* (generally 4 to 11 mg/ml) ***
Total Protein:	**Lot Specific***

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number:	**FM0075-FITC**
Clone:	**RB6-8C5**
Isotype:	Rat IgG2b, κ
Conjugation:	FITC
Excitation Source:	Blue 488 nm
Excitation Max:	494 nm
Emission Max:	518 nm
Immunogen:	Mouse granulocytes
Reported Applications:	Flow cytometry Immunohistochemistry (paraffin) Immunohistochemistry (frozen)
Formulation:	PBS, pH 7.0 Contains 0.09% Sodium Azide
Production:	Purified from cell culture supernatant in an animal-free facility
Purification:	Protein G
RRID:	AB_10312146

Description

The RB6-8C5 monoclonal antibody reacts strongly with mouse Ly6G and weakly with mouse Ly6C, previously referred to as GR-1. Ly6G is a 21-25 kDa member of the Ly-6 superfamily of GPI-anchored cell surface proteins with a role in cell signaling and cell adhesion. Ly6G is expressed differentially during development by cells in the myeloid lineage, including monocytes, macrophages, granulocytes, and neutrophils. Monocytes typically express Ly6G transiently during development, while mature granulocytes and peripheral neutrophils retain expression, making Ly6G a good cell surface marker for these populations. Studies have shown that the RB6-8C5 antibody inhibits the binding of the 1A8 antibody, which specifically reacts with mouse Ly6G but not Ly6C. This fluorescein isothiocyanate (FITC)-conjugated version of the RB6-8C5 antibody is useful for flow cytometry, immunofluorescence, immunohistochemistry (frozen), and immunohistochemistry (paraffin) applications.

Storage

Store at the stock concentration at 4°C and protected from prolonged exposure to light . **Do not freeze.**

Protocol Information

It is recommended that the reagent be carefully titrated for optimal performance in the assay of interest.

Application References

For a complete list of references, visit https://bioxcell.com/fm0075-fitc?bxcs=9k1b3a#tab_references or scan the QR code below.

Binding Validation

Validation data shown below confirms that this clone binds to its target antigen. For lot specific binding validation data, e-mail technicalservice@bioxcell.com.



Bio X Cell, LLC
https://bioxcell.com
+1-866-787-3444
customerservice@bioxcell.com