

Technical Data Sheet

ReadyTag anti-OVA



Attention: Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

Lot Specific Information

Lot Number: Lot Specific*
Volume: Lot Specific*
Concentration: Lot Specific* (generally 4 to 11 mg/ml) *
Total Protein: Lot Specific*

*This information will be noted on the certificate of analysis that ships with this product.

Product Information

Catalog Number: RT0267
Clone: F2-3.58
Isotype: Mouse IgG1
Recommended Dilution Buffer: InVivoPure pH 7.0 Dilution Buffer
Immunogen: Phosphoprotein constituent of egg white
Formulation: PBS, pH 7
Contains no stabilizers or preservatives
Endotoxin: <2EU/mg (<0.002EU/μg)
Determined by LAL gel clotting assay
Purity: >95%
Determined by SDS-PAGE
Sterility: 0.2 μm filtration
Production: Purified from cell culture supernatant in an animal-free facility
Purification: Protein G
RRID: [AB_2651130](https://eutils.ncbi.nlm.nih.gov/entrez/eutils/rrid.cgi?db=AB_2651130)
Molecular Weight: 150 kDa

Description

The F2-3.58 monoclonal antibody reacts with ovalbumin (OVA). OVA is a 45 kDa protein found in egg white that is commonly used in several different research areas. The protein is used to stimulate allergic inflammation in laboratory animals and is an established model allergen for airway hyper-responsiveness. OVA is also used in proteomics as a molecular weight marker.

Storage

Store at the stock concentration at 4°C. **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

Application References

For a complete list of references, visit https://bioxcell.com/rt0267?bxcs=9k1b3a#tab_references or scan the QR code below.



Bio X Cell, LLC

<https://bioxcell.com>

+1-866-787-3444

customerservice@bioxcell.com

Conditions: For research use only. Not for use in diagnostic or therapeutic procedures.

Not for resale.

Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC