

# Technical Data Sheet

## InVivoSIM anti-human TROP-2 (Sacituzumab Biosimilar)



**Attention:** Use of this product constitutes an agreement to Bio X Cell's Terms and Conditions which are included with this product in print and can also be found at <https://bioxcell.com/terms-and-conditions>.

### Lot Specific Information

**Lot Number:** Lot Specific\*  
**Volume:** Lot Specific\*  
**Concentration:** Lot Specific\* (generally 4 to 11 mg/ml) \*  
**Total Protein:** Lot Specific\*

\*This information will be noted on the certificate of analysis that ships with this product.

### Product Information

**Catalog Number:** SIM0051  
**Clone:** Sacituzumab  
**Isotype:** Human IgG1,  $\kappa$   
**Recommended Isotype Control(s):** RecombiMAb human IgG1 (K214R/L234F/L235E/P331S) isotype control, anti-hen egg lysozyme  
**Recommended Dilution Buffer:** InVivoPure pH 7.0 Dilution Buffer  
**Mutations:** K214R  
**Immunogen:** Human TROP-2  
**Reported Applications:** *in vitro* functional assays  
*in vivo* functional assays  
Antibody-drug conjugate synthesis  
ELISA  
Western blot  
Flow cytometry  
**Formulation:** PBS, pH 7.0  
Contains no stabilizers or preservatives  
**Endotoxin:** <0.5EU/mg (<0.0005EU/ $\mu$ g)  
Determined by LAL gel clotting assay  
**Purity:** >95%  
Determined by SDS-PAGE  
**Sterility:** 0.2  $\mu$ m filtration  
**Production:** Purified from cell culture supernatant in an animal-free facility  
**Aggregation:** <5%  
Determined by SEC  
**RRID:**  
**Molecular Weight:** 150 kDa

### Murine Pathogen Test Results

Mouse Norovirus: Negative, Mouse Parvovirus: Negative, Mouse Minute Virus: Negative, Mouse Hepatitis Virus: Negative, Reovirus Screen: Negative, Lymphocytic Choriomeningitis virus: Negative, Lactate Dehydrogenase-Elevating Virus: Negative, Mouse Rotavirus: Negative, Theiler's Murine Encephalomyelitis: Negative, Ectromelia/Mousepox Virus: Negative, Hantavirus: Negative, Polyoma Virus: Negative, Mouse Adenovirus: Negative, Sendai Virus: Negative, Mycoplasma Pulmonis: Negative, Pneumonia Virus of Mice: Negative, Mouse Cytomegalovirus: Negative, K Virus: Negative

### Description

This biosimilar antibody uses the same variable regions as the therapeutic antibody Sacituzumab, making it ideal for

research use. Sacituzumab is a fully humanized IgG1k monoclonal antibody that reacts with human trophoblastic cell surface antigen 2 (TROP-2), an antigen linked to cancer. TROP-2 is a cell surface receptor that spans the cellular membrane with an extracellular, transmembrane, and intracellular domain, along with a cytoplasmic tail essential for signaling. The ligands of TROP-2 include claudin-1, claudin-7, cyclin D1, and potentially IGF-1. TROP-2 upregulates EpCAM-triggered cell signaling and requires RIP to regulate efficient growth and division of cancer cells. Several proteins, including RIP, TACE,  $\gamma$ -secretase, Presenilin 1 (PS-1) and PS-2, facilitate TROP-2's cleavage within the transmembrane domain, and this cleavage produces TROP-2 extracellular domain (ECD) and intracellular domain (ICD) fragments. TROP-2 is considered a cellular marker of trophoblastic stem cells, and this glycoprotein is known to transduce calcium signals. Normal epithelial cells express TROP-2 at a low level, while many tumors (such as glioblastoma, pancreatic carcinoma, and all breast cancer subtypes, especially triple-negative breast cancer) express it at a high level. The critical role of TROP-2 in multiple signaling pathways often links to the proliferation, invasion, and metastasis of tumors. The tumor cell-specific expression of TROP-2 makes it an ideal target for cancer immunotherapy. Several recent studies have used Sacituzumab-based antibody-drug conjugates (ADC), namely Sacituzumab Govitecan (IMMU-132) that contains Sacituzumab conjugated to SN-38 (the active metabolite of irinotecan), utilizing a pH hydrolysable linker CL2A for facilitating SN-38 release to the cancer cells in in vitro and in vivo preclinical studies.

## Storage

Store at the stock concentration at 4°C . **Do not freeze.**

It is not uncommon for a floccule or precipitate to appear during storage. The floccule is typically buffer salts precipitating out of solution or a small bit of protein aggregation. For information on how to remove floccules or precipitates see our FAQ's at <https://bioxcell.com/faqs>.

## Protocol Information

Since applications vary, each investigator should use the application references as a guide to help estimate the appropriate dose or concentration. The dose or concentration can be further optimized experimentally in a dose response or titration experiment.

## Application References

For a complete list of references, visit [https://bioxcell.com/sim0051?bxcs=9k1b3a#tab\\_references](https://bioxcell.com/sim0051?bxcs=9k1b3a#tab_references) or scan the QR code below.



---

**Bio X Cell, LLC**

<https://bioxcell.com>

+1-866-787-3444

[customerservice@bioxcell.com](mailto:customerservice@bioxcell.com)

*Conditions: For research use only. Not for human use. Not for use in diagnostic or therapeutic procedures.*

*Not for resale.*

**Bio X Cell, Bio X Cell logo, and all other trademarks are the property of Bio X Cell, LLC © 2024 Bio X Cell, LLC**